

PHYSIOLOGICAL PRINCIPLES UNDERLYING SYNCHRONIZATION OF ESTRUS

M.F. Smith, G.A. Perry, J.A. Atkins, E.M. Jinks, K.G. Pohler, and D.J. Patterson

Division of Animal Sciences, University of Missouri,
Columbia
Department of Animal and Range Sciences, South
Dakota State University


Overview

- Physiology of the bovine estrous cycle
- Hormones utilized for estrus synchronization
- Estrus synchronization products
- Hormonal management of the luteal phase
- Hormonal management of follicular waves
- Management considerations


Characteristics of the Estrous Cycle

- Length of the estrous cycle
 - Average 21 days (range 17 to 24 days)
 - Two follicular waves – 17 to 20 days
 - Three follicular waves – 21 to 24 days
- Estrus (standing heat)
 - 12 to 18 hours (range 8 to 30 hours)
- Ovulation
 - Approximately 30 hours after the beginning of standing estrus


Duration of Estrus




Influence of confinement area on the number of mounts per estrus ($P > 0.5$)






Influence of the number of cows in estrus on number of times cows are mounted in drylot or pasture





Ovarian Structures


- Graffian follicle
 
- Corpus luteum
 



Stages of the Estrous Cycle


- Follicular Phase
- Estrus
- Luteal Phase

Follicular and Estrus Phases




- Maturation of the preovulatory follicle
- Estrous behavior
- Ovulation (induced by the LH surge)

Luteal Phase




- P₄ = Progesterone – CIDR and MGA
- PGF = Prostaglandin F – Lutalyse, ProstaMate, In Synch, Estrumate, estroPLAN

Corpus Luteum and Pregnancy




- Maternal recognition of pregnancy – day 15 to 16
- Caution – PGF can induce abortion in pregnant cows


Physiology of Follicular Waves



Problem of Anestrus!



Problem of Anestrus!



Hormones utilized in Estrus Synchronization Protocols

- Progesterone/Progestins
- Prostaglandin F_{2α}
- GnRH

Progesterone

Biological Functions

- Inhibit estrus/ovulation
- Preparation for pregnancy
- Maintenance of pregnancy

Progestins/Progesterone

Role in Synchronization

- Inhibit estrus/ovulation
- Induce cyclicity
- Dominant follicle turnover

Prostaglandin F_{2α}

Biological Function

- Luteal regression in nonpregnant animals

Role in Synchronization

- Induce premature luteal regression

Gonadotropin Releasing Hormone (GnRH)

Biological Function

- Control secretion of LH
- Induces gonadotropin surge

Role in Synchronization

- Induce ovulation
- Synchronize follicular waves

Estrus Synchronization Products

Progestins

- Melengestrol Acetate
- EAZI-BREED CIDR

Prostaglandin F_{2α}

- Lutalyse
- ProstaMate
- In Synch
- Estrumate
- estroPLAN

Estrus Synchronization Products

GnRH

- Cystorelin
- Fertagyl
- Factryl
- OvaCyst

Hormonal management of the luteal phase


- Progestins
- Prostaglandin F_{2α}

Role of Progestins in Estrus Synchronization


- Synchronization of estrus in
 - beef heifers & cows
 - dairy heifers & cows
- Advances onset of puberty in heifers
- Advances return to estrus after calving in cows

Progestins


- Melengestrol Acetate - MGA
- Controlled Internal Drug Release - CIDR



Inhibition of Ovulation Following Long-term Progestin Treatment



Persistent Follicles




Prostaglandin F_{2α} (PGF)

(Lutalyse, ProstaMate, Estrumate, In Synch, estroPLAN)

- Causes CL regression
- No effect on noncycling cattle
- No induction of cyclicity
... No Jump-start
- Effective days 6 to 16 of the estrous cycle (day 0 = estrus)

PGF_{2α} – Effect of stage of the cycle



Hormonal management of follicular waves


- GnRH – Induces dominant follicle to ovulate
- Estradiol – dominant follicle turnover (↓FSH)
- Progesterone – dominant follicle turnover (↓LH)

GnRH


(Cystorelin, Factrel, Fertagyl, Ovacyst)

- Induces ovulation
- Synchronizes follicular waves
- Induces formation of a CL

GnRH – Induced Surge of LH



GnRH-Induced Ovulation During a Follicular Wave



Management considerations for selecting heifers and cows for synchronization of estrus

Before you start an estrous synch and AI program – Heifers

What has the pregnancy rate of your heifers been over the past few years?

Have your heifers received growth promoting implants?

Have you selected an appropriate target weight?

Heifers-contd.

Have your heifers attained 65% of their mature body weight?

What proportion of your heifers have a reproductive tract score of ≥ 4 ?

Before you start an estrous synch and AI program – Postpartum Cows

What has the pregnancy rate in your cows been over the past few years?

What is the current length of your breeding season?

Postpartum Cows – contd.

What proportion of your cows are cycling by the start of the breeding season?

What was the body condition score of your cows at calving?

Postpartum Cows – contd.

What is the current body condition score of your cows?

How many days postpartum will your cows be when estrus synchronization is initiated?

Postpartum Cows and Heifers

How much time can you devote to estrus detection?

Considering your handling facilities, how many cows can you breed in an hour?

